


Spontaneity of Redox Reactions

- Compare the reaction when zinc metal is placed in copper(II) solution and the reaction when copper metal is placed in a zinc solution

Demo: Comparing two Similar Redox Reactions



vs.



∴ $\text{Cu}^{2+}_{(aq)}$ must be a stronger O.A. than $\text{Zn}^{2+}_{(aq)}$
 ∴ $\text{Zn}_{(s)}$ must be a stronger R.A. than $\text{Cu}_{(s)}$

- Different atoms and ions have different abilities/strength to reduce or oxidize other atoms or ions
 - Atoms and ions are listed in tables, ranked according to their ability/tendency to oxidize or reduce other atoms or ions
- A table of oxidizing and reducing agents is arranged as follows:
 - A complete list is found on pg.7 of data booklet
 - Tables are usually always written as a list of reduction half-reactions in the forward direction

Table of Selected Standard Electrode Potentials*

Reduction Half-Reaction	Electrical Potential E° (V)
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-(\text{aq})$	+2.87
$\text{PbO}_2(\text{s}) + \text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{PbSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$	+1.69
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Au}(\text{s})$	+1.50
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.34
$\text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{SO}_3(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.17
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}(\text{aq})$	+0.15
$\text{S}(\text{s}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{aq})$	+0.14
$\text{AgBr}(\text{s}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s}) + \text{Br}^-(\text{aq})$	+0.07
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0.00
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}(\text{s})$	-0.14
$\text{AgI}(\text{s}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s}) + \text{I}^-(\text{aq})$	-0.15
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$	-0.26

most reactive/
strongest O.A.

become
weaker


become
weaker

most reactive/
strongest RA

in general



- We can use this table to predict if redox reactions will be spontaneous or not
- **Redox Spontaneity Rule:** a spontaneous redox reaction will occur only if the oxidizing agent (OA) is above the reducing agent (RA) in a redox table




EXAMPLES:

1. Would the reaction between silver ions and zinc metal be spontaneous?

Step 1: Identify which reactant is the oxidizing agent and which one is the reducing agent.


- Can use the table on page 7 of data book to see if the atom/ion is on the oxidizing agent or reducing agent side.
- Can use a net ionic equation to see which reactants are the reducing and oxidizing agents
- Can use oxidation numbers on both reactants and products. Therefore, you also need to be able to predict what the products would be if the reaction was spontaneous. Remember, metals form positive ions (and vice versa) and non-metals form negative ions (and vice-versa).
- Can use the general rule that the more positive reactant will be the oxidizing agent and the more negative reactant will be the reducing agent

Step 2: Located the oxidizing agent and the reducing agent on the table to determine the spontaneity of the reaction using the spontaneity rule.



$\text{Ag}^+(\text{aq})$ (O.A.) is higher on the table than $\text{Zn}(\text{s})$ (R.A.),
 \therefore spontaneous rxn!

2. Would a spontaneous reaction occur when tin (II) bromide solution is poured into acidified sodium dichromate solution?



since at least one O.A. ($Cr_2O_7^{2-}/H^+$) is higher than one R.A. ($Br^- \& Sn^{2+}$), the rxn is spontaneous!

Now try pg. 440 #5-8 blue box & #3,5-7 from review section & Practice Problems #1, 2


EXAMPLE: Use the following information to rank the oxidizing agents from strongest to weakest.

1. $2X_{(s)} + Y^{2+}_{(aq)} \rightarrow 2X^+ + Y_{(s)}$ $Y^{2+}_{(aq)}$ is higher than $X_{(s)}$
2. $L_{(s)} + X_{(s)} \rightarrow L^-_{(aq)} + X^+_{(aq)}$ $L_{(s)}$ is higher than $X_{(s)}$
3. $L^-_{(aq)} + Y^{2+}_{(aq)} \rightarrow \text{no reaction}$ $Y^{2+}_{(aq)}$ is lower than $L^-_{(aq)}$

Step 1: Look at one reaction at a time and identify the oxidizing agent and the reducing agent. Use the same four methods located on the previous page (table, oxidation numbers, net ionic equation, or general rule).

Step 2: Using the spontaneity rule, position the oxidizing agent relative to the reducing agent depending on whether the data indicated the reaction to be spontaneous or not. Remember a redox table always lists the oxidizing agents on the left side and the reducing agents on the right side.

Step 3: Repeat steps 1 & 2 for all other reactions given, but continue to position the oxidizing and reducing agents in one table



If the spontaneity of the reactions was arranged in a table format, what do you notice?

RA \ OA	$X^+_{(aq)}$	$L_{(s)}$	$Y^{2+}_{(aq)}$
$X_{(s)}$		✓	✓
$L^-_{(aq)}$	X		X
$Y_{(s)}$	X	✓	

← more "+" form
 strongest R.A. →
 weakest R.A. →
 b/c causes the least spontaneous rxns!
 ↑ more "-" form
 ↑ weakest O.A.
 ↑ Strongest O.A.
 b/c it causes the most spontaneous rxns


EXAMPLE: Using the following information, to develop a redox table and then answer the following questions.

- 1.) $B^+_{(aq)} + D_{(s)} \rightarrow D^+_{(aq)} + B_{(s)}$ $B^+_{(aq)}$ is higher than $D_{(s)}$
- 2.) $D_{(aq)} + C_{(s)} \rightarrow \text{No reaction}$ $D^+_{(aq)}$ is lower than $C_{(s)}$
- 3.) $C^+_{(aq)} + B_{(s)} \rightarrow B^+_{(aq)} + C_{(s)}$ $C^+_{(aq)}$ is higher than $B_{(s)}$

a. What is the strongest reducing and oxidizing agent?

S.R.A. is $D_{(s)}$
 S.O.A. is $C^+_{(aq)}$

b. What metal is the most reactive? $D_{(s)}$



Now try Practice Problems

Answers to Questions for Comprehension


Student Textbook page 440

- Q5.** (a) spontaneous
(b) nonspontaneous
(c) spontaneous
(d) nonspontaneous


Q6. Use (a) because the reaction will proceed spontaneously.

Q7. Any of the following metals will work: Fe, Zn, Cr, Al, Mg, Na, Ca, Ba, Li.


The reactions for Fe, Zn, Mg, Ca, and Ba, are similar:






The reactions for Al and Cr are similar



The reactions for Na and Li are also similar:






Q8. Any ion that is an oxidizing agent higher than tin(reducing agent), will be spontaneous. This could include Pb^{2+} , Fe^{3+} , Cu^{2+} , Cu^+ , Ag^+ , Au^{3+} , Co^{3+} , and etc. However to be a salt, the metal ions need to be in an ionic compound that is soluble in water. For example $\text{Pb}(\text{NO}_3)_2$, FeCl_3 , CuSO_4 , AgNO_3 , and so forth. The net ionic equations would be very similar (the anions are spectator ions, so they will not appear in the equation).



Section 12.1 Review Answers

Student Textbook page 440

3. Calcium ions will spontaneously react with Fe^{2+} , Ni^{2+} , and Ag^+ .



5. (a) The reaction will occur.

The complete balanced equation is: $2\text{AgNO}_3(\text{aq}) + \text{Cd}(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + \text{Cd}(\text{NO}_3)_2(\text{aq})$

The net ionic equation is: $2\text{Ag}^+(\text{aq}) + \text{Cd}(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + \text{Cd}^{2+}(\text{aq})$

The ionic equation is: $2\text{Ag}^+(\text{aq}) + 2\text{NO}_3^-(\text{aq}) + \text{Cd}(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + \text{Cd}^{2+}(\text{aq}) + 2\text{NO}_3^-(\text{aq})$

(b) A reaction will not occur.





(c) The reaction will occur.

The complete balanced reaction is: $2\text{Al}(\text{s}) + 3\text{HgCl}_2(\text{aq}) \rightarrow 2\text{AlCl}_3(\text{aq}) + 3\text{Hg}(\text{s})$

The net ionic equation is: $2\text{Al}(\text{s}) + 3\text{Hg}^{2+}(\text{aq}) \rightarrow 2\text{Al}^{3+}(\text{aq}) + 3\text{Hg}(\text{s})$

The ionic equation is: $2\text{Al}(\text{s}) + 3\text{Hg}^{2+}(\text{aq}) + 6\text{Cl}^-(\text{aq}) \rightarrow 2\text{Al}^{3+}(\text{aq}) + 3\text{Hg}(\text{s}) + 6\text{Cl}^-(\text{aq})$

6. Possible answers for each part are shown below, although many answers exist. In general, equations must be balanced, oxidizing agents must gain electrons, and reducing agents must lose electrons. Spontaneous reactions will only occur if the oxidizing agent is higher than the reducing agent (or the reducing agent is lower than the oxidizing agent).






7. (a) $\text{Na}(\text{l}) + \text{K}^+(\text{aq}) \rightarrow \text{K}(\text{l}) + \text{Na}^+(\text{aq})$





(b) The potassium ions are the oxidizing agents. Sodium is the reducing agent.

Practice Problems





1. Would a spontaneous reaction occur if iron (II) chloride ($\text{FeCl}_{2(\text{aq})}$) is poured into an aluminium can?
2. A student is required to store an aqueous solution of iron(III) nitrate. She has a choice of a copper, tin, iron, or silver container. Choose an appropriate container which would be most suitable for storing the solution.
3. Use the following spontaneous reactions to identify the strongest oxidizing agent and reducing agent.



4. Use the following reactions to identify the most reactive metal and the most reactive ion.



5. Four metals were placed into test tubes containing various ion solutions. Their resulting behaviour is shown by the equations below. Create a reduction half-reaction table according to the chemicals tendency to react. Identify the strongest oxidizing agent and reducing agent.



6. An analytical chemist reacts an unknown metal X with a copper(II) sulfate solution, plating out copper metal. Metal X does not react with aqueous zinc nitrate.
 - a. What is the order for these metal ions in decreasing tendency/ability to react?
 - b. What group of metals are eliminated as a possible identity of the unknown metal?
 - c. What other solutions might be chosen next to help identify the unknown metal?

